

GENERALIZATION OF THE GROSS-PERRY METRICS

M. Jakimowicz and J. Tafel

Institute of Theoretical Physics, University of Warsaw,
Hoża 69, 00-681 Warsaw, Poland, email: tafel@fuw.edu.pl

Abstract

A class of $SO(n+1)$ symmetric solutions of the $(N+n+1)$ -dimensional Einstein equations is found. It contains 5-dimensional metrics of Gross and Perry and Millward.

1 Introduction

An extra spatial dimension was introduced by Kaluza and Klein (see e.g. [1]) in order to unify electromagnetism and gravity. Recently, due to the string theory, extra dimensions became a permanent part of the high energy theoretical physics. For instance, in brane-world models (see [2] for a review) matter fields are confined to a four-dimensional brane and gravity can propagate in higher dimensional bulk. These higher dimensional models motivate studying Einstein's equations in $D > 4$ dimensions. Some techniques of the 4-dimensional Einstein theory were already generalized to higher dimensions. They refer mainly to the classification of the Weyl tensor, the Robinson-Trautman solutions, spacetimes with vanishing invariants and metrics with D-2 abelian symmetries (see e.g. [3, 4, 5, 6]).

Symmetry assumptions are one of the most efficient methods of solving the Einstein equations. All well known multidimensional exact solutions like the Myers-Perry black hole [7], black ring of Emparan and Reall [8] and the Gross-Perry metrics [9] (see also [10]) admit several dimensional symmetry groups.

In [11] we proposed a construction of vacuum metrics admitting $SO(n+1)$ spherical symmetry, which was based on the symmetry reduction of $(N+n+1)$ -dimensional Einstein equations to $(N+1)$ -dimensional equations with a scalar field ϕ . There was used an additional assumption that the field of normal vectors to surfaces $\phi = \text{const}$ is geodesic and the induced metric of the surfaces is an Einstein metric. The construction, for zero cosmological constant and timelike surfaces $\phi = \text{const}$, can be summarized as follows.

Let γ_{ij} and P_{ij} be symmetric tensors depending on coordinates x^i , $i = 0, \dots, N-1$. Assume that γ_{ij} has the Lorentz signature and P_{ij} satisfies the following conditions

$$P^i_i = 0, \tag{1}$$

$$P^i_j P^j_i = 2c = \text{const}, \tag{2}$$

$$P^k_{i;k} = 0, \tag{3}$$

where $P^i_j = \gamma^{ik} P_{kj}$ and the semicolon denotes covariant derivative related to the metric $\gamma_{ij} dx^i dx^j$. From matrices $\gamma = (\gamma_{ij})$ and $P = (P^i_j)$ we compose metric corresponding to $\gamma e^{P\tau}$, where τ is a function of another coordinate s . We assume that its Ricci tensor satisfies

$$R^i_j(\gamma e^{P\tau}) = \lambda \delta^i_j, \quad \lambda = \text{const}. \quad (4)$$

Given c and λ we look for solutions $\beta(s)$ and $\phi(s)$ of the following equations

$$(\beta \dot{\phi})^\cdot = -\beta V_{,\phi}, \quad (5)$$

$$-N\lambda\beta^{-2/N} = (1 - \frac{1}{N})\frac{\dot{\beta}^2}{\beta^2} - \frac{2c}{\beta^2} - \dot{\phi}^2 + 2V, \quad (6)$$

where the dot denotes the partial derivative with respect to s and V is a function of ϕ .

The main result of [11] is that, under conditions (1)-(6), metric

$$\tilde{g} = -ds^2 + \tilde{g}_{ij} dx^i dx^j, \quad (7)$$

were

$$\tilde{g}_{ij} = \beta^{2/N} (\gamma e^{P\tau})_{ij} \quad (8)$$

and $\tau(s)$ is defined via equation

$$\beta \dot{\tau} = 2, \quad (9)$$

satisfies $(N+1)$ -dimensional Einstein equations with the scalar field ϕ and potential V . Moreover, if

$$V = -\frac{1}{2}n(n-1)e^{-2\sqrt{\frac{n+N-1}{n(N-1)}}\phi} \quad (10)$$

then

$$g = e^{-2\sqrt{\frac{n}{(N-1)(n+N-1)}}\phi} \tilde{g} - e^{2\sqrt{\frac{N-1}{n(n+N-1)}}\phi} d\Omega_n^2 \quad (11)$$

is an $(N+n+1)$ -dimensional vacuum metric invariant under the group $SO(n+1)$. Here $d\Omega_n^2$ is the standard metric of the n -dimensional sphere.

A particular solution of conditions (1)-(4), for any $N > 1$, is given by

$$\gamma_{ij} = \text{diag}(+1, -1, -1, \dots), \quad P_{ij} = P_{ji} = \text{const}, \quad P^i_i = 0, \quad \lambda = 0. \quad (12)$$

For $N = 2$ conditions (1)-(4) can be solved in full generality. They lead either to (12) or to $c = 0$ and to $\gamma e^{P\tau}$ equivalent to the metric

$$(\gamma e^{P\tau})_{ij} dx^i dx^j = \frac{dudv}{(1 + \frac{\lambda}{4}uv)^2} + \tau h(u) du^2, \quad (13)$$

where h is an arbitrary function of coordinate u . In the next section we find solutions of equations (5), (6) and construct corresponding vacuum metrics. In section 3 we discuss properties of these metrics.

2 Multi dimensional vacuum metrics

In [11] we gave examples of vacuum metrics derived by our method. Other solutions with $n > 1$ can be obtained by inspection of the Gross-Perry metrics [9]. Let $\lambda = 0$ and $s = s(r)$ be a function of a new coordinate r . Then equations (5), (6), (9) take the form

$$\left(\frac{\beta\phi'}{\alpha}\right)' = \alpha\beta V_{,\phi} \quad (14)$$

$$\left(1 - \frac{1}{N}\right)\frac{\beta'^2}{\alpha^2} - \frac{\beta^2\phi'^2}{\alpha^2} + 2\beta^2 V = 2c, \quad (15)$$

$$\tau' = \frac{2\alpha}{\beta}, \quad (16)$$

where the prime denotes the derivative with respect to r and $\alpha = s'$. Metric (7) is given by

$$\tilde{g} = -\alpha^2 dr^2 + \beta^{2/N} (\gamma e^{P\tau})_{ij} dx^i dx^j. \quad (17)$$

If $N = n = 2$ equations (14), (15) are satisfied by functions α , β , ϕ corresponding to the Gross-Perry metric [9]. Changing parameters in these functions leads to the following solutions for arbitrary dimensions $N > 1$ and $n > 1$

$$\alpha = \alpha_0 |r|^{-l-1} |r - r_0|^{l-p} |r + r_0|^{l+p} \quad (18)$$

$$\beta = \beta_0 (r^2 - r_0^2) \alpha \quad (19)$$

$$e^{\sqrt{\frac{n+N-1}{n(N-1)}}\phi} = (n-1) |r\alpha|. \quad (20)$$

Here l is a number defined by n and N

$$l = \frac{n + N - 1}{(n - 1)(N - 1)} \quad (21)$$

and p , α_0 , β_0 and $r_0 \neq 0$ are parameters related to the constant c via

$$c = 2\beta_0^2 r_0^2 \left[\frac{n}{n-1} - p^2 \frac{(n-1)(N-1)^2}{N(n+N-1)} \right]. \quad (22)$$

Integrating equation (16) yields

$$\tau = \frac{1}{\beta_0 r_0} \ln \left| \frac{r + r_0}{r - r_0} \right| + \tau_0. \quad (23)$$

Due to a freedom of transformations of r , P and γ we can assume

$$r_0 > 0, \quad |\alpha_0| = \frac{1}{n-1}, \quad \beta_0 = 1, \quad \tau_0 = 0 \quad (24)$$

(note that a sign of α_0 can be still adjusted to have $\beta > 0$ for $r \neq 0, \pm r_0$). Thus, p and $r_0 > 0$ remain as free parameters.

Let $N = 2$. In the case (12) and $c > 0$ the matrix P can be diagonalized by a 2-dimensional Lorentz transformation. Hence, one obtains

$$\tilde{g} = -\alpha^2 dr^2 + \beta \left(e^{\pm \tau \sqrt{c}} dt^2 - e^{\mp \tau \sqrt{c}} dy^2 \right) , \quad (25)$$

where t and y denote coordinates x^i . Substituting (18)-(24) into (25) and (11) yields the following $(n+3)$ -dimensional vacuum metric

$$g = \left| \frac{r - r_0}{r + r_0} \right|^{p' - q} dt^2 - \left| \frac{r - r_0}{r + r_0} \right|^{p' + q} dy^2 - \frac{|r + r_0|^{\frac{2p' + 2}{n-1}}}{|r|^{\frac{2n}{n-1}} |r - r_0|^{\frac{2p' - 2}{n-1}}} \left(\frac{dr^2}{(n-1)^2} + r^2 d\Omega_n^2 \right) . \quad (26)$$

Parameters p' and q are related to p and c by

$$p' = \frac{n-1}{n+1} p , \quad q = \pm \frac{\sqrt{|c|}}{r_0} . \quad (27)$$

Because of (22) they are constrained by

$$(n+1)p'^2 + (n-1)q^2 = 2n . \quad (28)$$

For $n=2$ solution (26) is exactly the Gross-Perry metric [9] under the identification

$$r_0 = m, \quad p' = \frac{1}{\alpha}(\beta + 1), \quad q = \frac{1}{\alpha}(\beta - 1). \quad (29)$$

Here m , α and β are parameters used by Gross and Perry, constrained by the condition $\alpha = \sqrt{\beta^2 + \beta + 1}$.

If $c < 0$ the matrix P can be put into the off diagonal form. Instead of (25) one obtains

$$\tilde{g} = -\alpha^2 dr^2 + \beta \left[\cos(\tau \sqrt{|c|})(dt^2 - dy^2) \pm 2 \sin(\tau \sqrt{|c|}) dt dy \right] . \quad (30)$$

In this case the vacuum metric corresponding to (18)-(24) reads

$$g = \left| \frac{r - r_0}{r + r_0} \right|^{p'} \left[\cos \left(q \ln \left| \frac{r + r_0}{r - r_0} \right| \right) (dt^2 - dy^2) + 2 \sin \left(q \ln \left| \frac{r + r_0}{r - r_0} \right| \right) dt dy \right] - \frac{|r + r_0|^{\frac{2p' + 2}{n-1}}}{|r|^{\frac{2n}{n-1}} |r - r_0|^{\frac{2p' - 2}{n-1}}} \left(\frac{dr^2}{(n-1)^2} + r^2 d\Omega_n^2 \right) . \quad (31)$$

Relation (27) is still valid, but now parameters p' , q are constrained by

$$(n+1)p'^2 - (n-1)q^2 = 2n . \quad (32)$$

If $N = 2$ and

$$p = \pm \frac{\sqrt{2n(n+1)}}{n-1} \quad (33)$$

then it follows from (22) that $c = 0$ and one can merge solutions (18)-(20) with metric (13) for $\lambda = 0$. In this way the following vacuum metric is obtained

$$g = \left| \frac{r-r_0}{r+r_0} \right|^{\pm \sqrt{\frac{2n}{n+1}}} \left(dudv + \ln \left| \frac{r+r_0}{r-r_0} \right| h(u) du^2 \right) - \frac{|r+r_0|^{\frac{2}{n-1}(\pm \sqrt{\frac{2n}{n+1}}+1)}}{|r|^{\frac{2n}{n-1}}|r-r_0|^{\frac{2}{n-1}(\pm \sqrt{\frac{2n}{n+1}}-1)}} \left(\frac{dr^2}{(n-1)^2} + r^2 d\Omega_n^2 \right). \quad (34)$$

In the case $n = 2$, $h(u) = 0$ metric (34) with the lower sign coincides with the metric given by Millward [12] under the identification

$$b = \frac{1}{\sqrt{3}} \ln \left| \frac{r-r_0}{r+r_0} \right|, \quad M = \frac{\sqrt{3}}{2} r_0. \quad (35)$$

For $N > 2$ one can easily construct vacuum solutions based on relations (11), (12), (17)-(24). They generalize metrics (26) and (31). In this case a classification of symmetric tensors (here P_{ij}) in multidimensional Lorentzian manifolds [3] can be useful in order to distinguish nonequivalent solutions. One can also construct metrics which generalize (34) by taking $\gamma e^{P\tau}$ corresponding to the metric

$$dudv + \tau h(u) du^2 + \sum_{a=1}^{N-2} e^{c_a \tau} dy_a^2, \quad (36)$$

where constants c_a are constrained by

$$\sum_{a=1}^{N-2} c_a = 0. \quad (37)$$

In this case we can use functions defined by (18)-(24) with constant c given by

$$c = \frac{1}{2} \sum_{a=1}^{N-2} c_a^2. \quad (38)$$

3 Discussion

In addition to $SO(n+1)$ symmetries metrics (26) and (31) admit one timelike and one spacelike Killing vector (note that interpretation of ∂_t and ∂_y in case (31) can change depending on value of r). Metric (26) is static and metric (31) is stationary. In the limit $r \rightarrow \infty$ they behave like

$$dt^2 - dy^2 - r^{-2\frac{n-2}{n-1}} \left(\frac{dr^2}{(n-1)^2} + r^2 d\Omega_n^2 \right). \quad (39)$$

Under the change $r' = r^{\frac{1}{n-1}}$ metric (39) takes the standard form of the (n+3)-dimensional Minkowski metric. Thus, metrics (26) and (31) are asymptotically flat on surfaces $y = \text{const.}$

Metric (34) has a null Killing vector field ∂_v and it belongs to generalized Kundt's class [14]. If $r \rightarrow \infty$ it tends asymptotically to the flat metric in the form

$$dudv - r^{-2\frac{n-2}{n-1}} \left(\frac{dr^2}{(n-1)^2} + r^2 d\Omega_n^2 \right). \quad (40)$$

Generalizing results of [13] for the Gross-Perry metric to arbitrary n, one can show that both metrics (26) and (31) are of algebraic type *I*. For $h \neq 0$ metric (34) is of algebraic type *II*_i and for $h(u) = 0$ it is of type *D*. Aligned null vector fields for metrics (26), (31) and (34) are given in Appendix A.

All metrics (26), (31) and (34) are singular at $r = \pm r_0$ and $r = 0$. Near $r = 0$ they behave as

$$dt^2 - dy^2 - r^{-\frac{2n}{n-1}} \left(\frac{dr^2}{(n-1)^2} + r^2 d\Omega_n^2 \right). \quad (41)$$

Substituting $r' = r^{-\frac{1}{n-1}}$ shows that (41) is the flat metric. Thus, $r = 0$ is a coordinate singularity. By calculating the Kretschmann invariant (see Appendix B) it can be shown, that singularity at $r = \pm r_0$ is essential for all values of parameters in the case of metrics (26) and (34). In the case of metric (31) the singularity at $r = r_0$ and $r = -r_0$ is essential when, respectively, $p' < n$ or $p' > -n$. For $p' > n$ or $p' < -n$ the geodesic distance along ∂_r tends to infinity when $r \rightarrow r_0$ or $r \rightarrow -r_0$, respectively. Thus, these regions represents an infinity different from that given by $r \rightarrow \infty$. For these values of parameters the Riemann tensor (in an orthonormal basis) tends to zero when $r \rightarrow r_0$ or $r \rightarrow -r_0$, respectively. However, the asymptotic metric is not the (n+3)-dimensional Minkowski metric. Its coefficients in front of dt and dy tend to zero whereas the coefficient in front of $d\Omega_n^2$ tends to infinity like the geodesic distance to the power $2(p' - 1)/(p' - n)$ or $2(p' + 1)/(p' + n)$, respectively.

Since metrics (26), (31) are invariant under the nonnull field ∂_u they can be interpreted in the context of the Kaluza-Klein theory. Then metric (26) is equivalent to the scalar field given by g_{yy} and the asymptotically flat (n+2)-dimensional metric induced on the surface $y=\text{const.}$ The case $n = 2$ (the Gross-Perry metric) was studied in this framework by Ponce de Leon [15]. In order to interpret metric (31) with $q \neq 0$ in this vein one can write it in the form

$$g = -\Phi(dy - A_0 dt)^2 + g_{n+2}. \quad (42)$$

Here

$$A_0 = \tan \left(q \ln \left| \frac{r + r_0}{r - r_0} \right| \right) \quad (43)$$

is the electromagnetic potential,

$$\Phi = \left| \frac{r - r_0}{r + r_0} \right|^{p'} \cos \left(q \ln \left| \frac{r + r_0}{r - r_0} \right| \right) \quad (44)$$

corresponds to a scalar field and

$$g_{n+2} = \left| \frac{r-r_0}{r+r_0} \right|^{p'} \frac{dt^2}{\cos \left(q \ln \left| \frac{r+r_0}{r-r_0} \right| \right)} - \frac{|r+r_0|^{\frac{2p'+2}{n-1}}}{|r|^{\frac{2n}{n-1}} |r-r_0|^{\frac{2p'-2}{n-1}}} \left(\frac{dr^2}{(n-1)^2} + r^2 d\Omega_n^2 \right) \quad (45)$$

defines, modulo a power of Φ , a $(n+2)$ -dimensional metric. This metric is Lorentzian and asymptotically flat for large values of r and becomes singular when r diminishes to a value satisfying condition $q \ln \left| \frac{r+r_0}{r-r_0} \right| = \pm \pi/2$.

Acknowledgements. This work was partially supported by the Polish Committee for Scientific Research (grant 1 PO3B 075 29).

Appendix A

The aligned null direction is given by

$$\begin{aligned} \hat{l} = & \left(\left| \frac{r-r_0}{r+r_0} \right|^{p'-q} + \frac{1}{2} f^2 \right) dt + \left(\left| \frac{r-r_0}{r+r_0} \right|^{p'} - \frac{1}{2} f^2 \left| \frac{r-r_0}{r+r_0} \right|^q \right) dy + \\ & + f \frac{|r+r_0|^{\frac{p'+1}{n-1}}}{(n-1) |r|^{\frac{n}{n-1}} |r-r_0|^{\frac{p'-1}{n-1}}} dr \end{aligned} \quad (A.1)$$

for metric (26) and by

$$\begin{aligned} \hat{l} = & \left(\sin \left(q \ln \left| \frac{r+r_0}{r-r_0} \right| \right) + 1 \right) \left(1 - \frac{1}{2} f^2 \left| \frac{r-r_0}{r+r_0} \right|^{p'} \frac{\cos \left(q \ln \left| \frac{r+r_0}{r-r_0} \right| \right)}{\left(\sin \left(q \ln \left| \frac{r+r_0}{r-r_0} \right| \right) + 1 \right)^2} \right) dt + \\ & + \left(\frac{1}{2} f^2 \left| \frac{r-r_0}{r+r_0} \right|^{p'} - \cos \left(q \ln \left| \frac{r+r_0}{r-r_0} \right| \right) \right) dy + f \frac{|r+r_0|^{\frac{p'+1}{n-1}}}{(n-1) |r|^{\frac{n}{n-1}} |r-r_0|^{\frac{p'-1}{n-1}}} dr \end{aligned} \quad (A.2)$$

for metric (31). The function f is a solution of the polynomial equation

$$f^4 - 8f^2 \left(\frac{p'}{q} + \frac{2q(n-1)rr_0}{n(r^2 + r_0^2 - 2p'rr_0)} \right) A - 16A^2 = 0, \quad (A.3)$$

where

$$A = \left| \frac{r-r_0}{r+r_0} \right|^{p'-q} \quad (A.4)$$

in the case (26) and

$$A = \left| \frac{r-r_0}{r+r_0} \right|^{-p'} \left(\sin \left(q \ln \left| \frac{r+r_0}{r-r_0} \right| \right) + 1 \right) \quad (A.5)$$

in the case (31). For metric (34) the aligned null directions are defined by

$$\hat{n} = \left| \frac{r + r_0}{r - r_0} \right|^{\pm \sqrt{\frac{2n}{n+1}}} du \quad (\text{A.6})$$

and

$$\hat{l} = \frac{1}{2} \left(\ln \left| \frac{r - r_0}{r + r_0} \right| + f^2 \left| \frac{r + r_0}{r - r_0} \right|^{\pm \sqrt{\frac{2n}{n+1}}} \right) du + \frac{1}{2} dv + \frac{|r + r_0|^{\frac{1}{n-1}(\pm \sqrt{\frac{2n}{n+1}} + 1)}}{(n-1)|r|^{\frac{n}{n-1}}|r - r_0|^{\frac{1}{n-1}(\pm \sqrt{\frac{2n}{n+1}} - 1)}} dy. \quad (\text{A.7})$$

In this case

$$f^2 = \left(\mp \sqrt{\frac{n+1}{2n}} + \frac{2(n^2 - 1)rr_0}{n(n+1)(r^2 + r_0^2) \mp \sqrt{2n(n+1)}rr_0} \right) h(u) \left| \frac{r + r_0}{r - r_0} \right|^{\mp \sqrt{\frac{2n}{n+1}}}. \quad (\text{A.8})$$

Appendix B

The Kretschmann invariant for metrics (26) and (31) has the following form

$$\begin{aligned} R_{\mu\nu\delta\sigma}R^{\mu\nu\delta\sigma} = & 16n(n-1)r_0^2|r|^{\frac{2(n+1)}{n-1}}|r-r_0|^{-\frac{4(n-p')}{n-1}}|r+r_0|^{-\frac{4(n+p')}{n-1}} \left((n^2+n-2p'^2)r^4 + \right. \\ & + 2p'(n(n+1)(p'^2-3) + 2(1+p'^2))r^3r_0 + \\ & - (4-3n-5n^2-2(-2+n(n+3))p'^2 + (4+n(n+3))p'^4)r^2r_0^2 + \\ & \left. + 2p'(n(n+1)(p'^2-3) + 2(1+p'^2))rr_0^3 + (n^2+n-2p'^2)r_0^4 \right). \quad (\text{B.1}) \end{aligned}$$

In the case of metric (34) the Kretschmann invariant is given by (B.1) with $p' = \pm \sqrt{\frac{2n}{n+1}}$.

References

- [1] Coquereaux R. and Jadczyk A. 1988, Riemannian geometry, fiber bundles, Kaluza-Klein theories and all that..., *World Scientific Lecture Notes in Physics* **16**
- [2] Maartens R. 2004, Brane-world gravity, *Living Rev. Relativity* **7**, (<http://www.livingreviews.org/lrr-2004-7>)
- [3] Coley A., Milson R., Pravda V., Pravdova A. 2004, Classification of the Weyl Tensor in Higher Dimensions *Class. Quantum Grav.* **21** L35-L42
- [4] Podolsky J. and Ortaggio M. 2006, Robinson-Trautman spacetimes in higher dimensions, *Class. Quantum Grav.*, **23** 57855797
- [5] Coley A., Fuster A., Hervik S., Pelavas N. 2006, Higher dimensional VSI spacetimes, *Class. Quantum Grav.*, **23** 7431-7444

- [6] Harmark T. 2004, Stationary and Axisymmetric Solutions of Higher-Dimensional General Relativity, *Phys. Rev.*, **D70** 124002
- [7] Myers R.C. and Perry M.J. 1986, Black holes in higher dimensional space-time, *Annals Phys.*, **172** 304
- [8] Emparan R. and Reall H.S. 2002, Rotating black ring in five dimensions, *Phys. Rev. Lett.*, **88** 101101
- [9] Gross D.J. and Perry M.J. 1983, Magnetic monopoles in Kaluza-Klein theories, *Nucl. Phys.*, **B 226** 29
- [10] Davidson A. and Owen D. 1985, Black holes as a windows to extra dimensions *Phys. Lett.*, **B 155** 247
- [11] Jakimowicz M. and Tafel J. 2008, $SO(n + 1)$ symmetric solutions of the Einstein equations in higher dimensions, *Class. Quantum Grav.*, **25** 175002
- [12] Millward R.S. 2008, A five-dimensional Schwarzschild-like solution, arXiv: gr-qc/0603132
- [13] Coley A. and Pelavas N. 2006, Classification of Higher Dimensional Spacetimes, *Gen. Rel. Grav.* **38** 445-461
- [14] Coley A. 2008, Classification of the Weyl Tensor in Higher Dimensions and Applications, *Class. Quantum Grav.*, **25** 033001
- [15] Ponce de Leon J. 2007, Exterior spacetime for stellar models in 5-dimensional Kaluza-Klein gravity, *Class. Quantum Grav.* **24** 1755